


Lampiran 1. Skema Kerja Penelitian



Lampiran 2. Buah mengkudu (*Morinda citrifolia* L)



Gambar 1. Mengkudu segar




Gambar 2. Mengkudu mengkal yang sudah diiris



Gambar 3. Mengkudu kering


Lampiran 3. Diameter zona bening ekstrak mengkudu (*Morinda citrifolia* L) terhadap *Staphylococcus aureus*

Ekstrak	I	II	III	Rata-rata
Heksan	1,2 cm	1,0 cm	0,0 cm	0,7 cm
Etil asetat	0,0 cm	0,0 cm	0,0 cm	0,0 cm
Butanol	1,2 cm	0,0 cm	0,0 cm	0,4 cm
Jus	2,3 cm	2,2 cm	2,5 cm	2,3 cm
Kontrol (+)	2,8 cm	3,0 cm	2,9 cm	2,9 cm




Lampiran 4. Diameter zona bening ekstrak mengkudu (*Morinda citrifolia* L) terhadap *Candida albicans*

Uji	I	II	III	Rata-rata
Ekstrak Heksan 1%	1,1 cm	1 cm	1,1 cm	1,07 cm
Ekstrak Etis Asetat 1%	1 cm	0	0,9 cm	0,63 cm
Ekstrak Butanol 1%	0,8 cm	0	0	0,27 cm
Jus 1%	0	0	1 cm	0,33 cm
Etanol	0	0	0	0
Akuades	0	0	0	0
Ketokonazol 1%	1,2 cm	1,3 cm	1,1 cm	1,2 cm




Lampiran 5. Diameter zona bening jus mengkudu matang (*Morinda citrifolia* L) terhadap *Staphylococcus aureus*

Jus	I	II	III	Rata-rata
1%	2 cm	1,6 cm	2,1 cm	1,9 cm
3%	1,2 cm	1 cm	1,6 cm	1,3 cm
5%	1,4 cm	0,7 cm	0,8 cm	1,0 cm
7%	1 cm	0,7 cm	0,7 cm	0,8 cm
Kontrol (+)	3 cm	2,9 cm	3 cm	3,0 cm



**Lampiran 6. Diameter zona bening jus mengkudu matang (*Morinda citrifolia* L.)
terhadap *Candida albicans***


Uji	I	II	III	Rata-rata
Jus 1%	0,8 cm	0	0	0,27 cm
Jus 3%	0,9 cm	0	0	0,30 cm
Jus 5%	1 cm	0	0	0,33 cm
Jus 7%	1 cm	0	0	0,33 cm
Ketokonazol 1%	1,2 cm	1,1 cm	1,4 cm	1,23 cm
Etanol	0	0	0	0
Akuades	0	0	0	0



Lampiran 7. Analisis aktivitas antibakteri salep terhadap *Staphylococcus aureus*

Gambar 1. Dasar salep F1a

Gambar 1. F1a dan F1b




Gambar 2. Dasar salep F2a

Gambar 2. F2a dan F2b



Gambar 3. F3a dan F3b



Gambar 4. F4a dan F4b



Gambar 5. Salep aktif F1b



Gambar 6. Salep aktif F4b

